

SUGARCANE (*SACCHARUM OFFICINARUM* L) RESPONSE TO INTEGRATED APPLICATION OF INORGANIC AND ORGANIC FERTILIZERS FOR YIELD AND QUALITY ATTRIBUTES

Dr. Abdul Fatah Soomro and Riaz Noor Panhwar

National Sugar and Tropical Horticulture Research Institute, PARC, Makli, Thatta

ABSTRACT

To study the response of sugarcane to inorganic fertilizer application integrated with organic nutrient sources for cane yield and quality parameters, field experiment was carry out at Agriculture Research Institute, Tandojam, Pakistan, (25°25'60"N 68°31' 60E) during 2008-2009. Sugarcane variety Thatta-10 was planted under RCBD. The treatments combinations were as: control (0-0-0), Recommended NPK dose (225-112-168), FYM 10 t ha⁻¹, FYM 20 t ha⁻¹, PM 10 t ha⁻¹, PM 20 t ha⁻¹, BF 5 t ha⁻¹, BF 10 t ha⁻¹, three-fourth of recommended rate (169-84-126)+ FYM 10 t ha⁻¹, three-fourth of recommended rate (169-84-126) + FYM 20 t ha⁻¹, three-fourth of recommended rate (169-84-126) + PM 10 t ha⁻¹, three-fourth of recommended rate (169-84-126) + PM 20 t ha⁻¹, three-fourth of recommended rate (169-84-126) + BF 5 t ha⁻¹, three-fourth of recommended rate (169-84-126) + BF 10 t ha⁻¹ , half of recommended rate(112-56-84) + FYM 10 t ha⁻¹, half of recommended rate(112-56-84) + FYM 20 t ha⁻¹, half of recommended rate (112-56-84) + PM 10 t ha⁻¹, half of recommended rate(112-56-84) + PM 20 t ha⁻¹, half of recommended rate(112-56-84)+BF 5t ha⁻¹, and half of recommended rate(112-56-84) + BF 10 t ha⁻¹. The results showed that sole application of inorganic or organic nutrient sources exhibited no prominent increase in all sugarcane traits but the integration of both the sources except bio fertilizer significantly improved all the traits of plant crop. The maximum plant height, stem girth, , millable canes, cane yield, leaf area plant⁻¹, leaf area index, crop growth rate and dry matter were found higher with the application of FYM and or press mud applied at 20 t ha⁻¹ with three-fourth of recommended rate of NPK fertilizer (169-84-126). Both the treatments were statistically non significant. However, quality and nutrient uptake traits viz. brix, pol, purity, commercial cane sugar, NPK uptake and accumulation in sugarcane were higher with the application of three-fourth of recommended rate of NPK fertilizer (169-84-1261) + 20 tons press mud ha⁻¹. It is concluded that integrated nutrient management showed 25% saving of inorganic fertilizers with the application of FYM and or press mud applied at 20 t ha⁻¹. Integration of organic and inorganic nutrients should be practiced. This will not only enhance growth, yield, quality and nutrient uptake of sugarcane but also conserve agro-ecosystem for sustainable crop production.

INTRODUCTION

Sugarcane (or *Saccharum hybrid*) is an important cash crop in Pakistan. It provides sucrose, alcohol and organic matter waste which is utilized as fertilizer. Sugarcane was cultivated vegetatively as noble canes until the end of 20th century. Currently sugarcane is recognized as an aneuploid hybrid crop obtained from a cross between *Saccharum officinarum*

L., and a wild relative *Saccharum spontaneum*, through the process of Nabilization (Mumtaz *et al.*, 2011). The use of inorganic fertilizers does not necessarily lead to better farming than the use of natural and organic methods in agriculture. Due to continuous application of only inorganic fertilizers and plant protection chemicals in agriculture, the soils have been badly degraded. It has destroyed stable traditional ecosystem of the soil (Palaniappan & Annadurai, 1999). There is need to encourage more productive, cost efficient and ecofriendly farming system (Bhattacharya & Gehlot, 2003). The use of organic manure has been the need for improving the sustainable productivity of soil. (Singh *et al.*, 2003) depicted that by using integration of 25 and 37.5% N through sulphitation press mud cake and rest through inorganic nitrogen, the inorganic N could be saved from 12.5 to 25% with no loss in cane yield as obtained against the total N application (150 Kg/ha) as inorganic source . However, 50% N through SPMC + 50% through inorganic sources increased the cane yield significantly being 12.05% higher than that of treatment fertilized with 150 kg N/ha as inorganic source alone. Like micronutrients, farm yard manure (FYM) is also considered as an important source of macro and micronutrients that increase crop yield. Due to higher prices of inorganic fertilizers, farmers in Pakistan could easily manage to prepare FYM in their farms and to apply in fields Thus use of Farmyard Manure, Potassium and Zinc should be included in integrated crop management approaches for sustainable agriculture (Nawab *et al.*, 2011). Addition of compost improves soil structure, texture and tilth. Biocomposts have gained importance since the fertilizers and pesticides cause a lot of environmental problems and health hazards and soil degradation (Ghugare *et al.*, 1988). Organic matter is necessary for sustainable crop production (Bhander *et al.*, 1998). Sugarcane is a long duration and exhaustive crop, which produces large quantum biomass, removes considerable amount of nutrients from soil for its normal growth and development. A crop of 100 ton cane yield may remove 140 kg N, 34 kg P and 332 kg K from soil (Dang *et al.*, 1995). Comparisons of multiple organic and inorganic sources are useful nutrient management options that can improve sugarcane yields. Increasing nutrient management options are particularly important during increasing fertilizer prices. Intensive cropping and imbalanced use of essential plant nutrients have rendered the alluvial soils of sub-tropics to be poor in organic carbon content and deterioration in physical properties (Speir *et al.*, 2004) lead to restricted growth and development of the crop. For sustainable crop production, integrated use of chemical and organic fertilizer has proved to be highly beneficial. Several researchers have demonstrated the beneficial effect of combined use of

chemical and organic fertilizers to mitigate the deficiency of many secondary and micronutrients in fields that continuously received only N, P and K fertilizers for a few years, without any micronutrient or organic fertilizer. Dutta *et al.*, (2003) reported that the use of organic fertilizers together with chemical fertilizers, compared to the addition of organic fertilizers alone, had a higher positive effect on microbial biomass and hence soil health. However application of organic manure in combination with chemical fertilizer has been reported to increase absorption of N, P and K in sugarcane leaf tissue in the plant and ratoon crop, compared to chemical fertilizer alone (Bokhtiar & Sakurai 2005). Application of organic fertilizers together with chemical fertilizers, compared to the addition of organic fertilizers alone, had a higher positive effect on microbial biomass and hence soil health (Kumaraswamy *et al.*, 1998). The application of organic matter from such resources as animal manure, crop residues and green manuring has been shown to replenish soil organic C and improve soil fertility (Saviozzi *et al.*, 2002; Srivastava *et al.*, 2009). Moreover several kind of microbial agents capable of fixing N or mobilizing P and others nutrients are becoming an integral component of Integrated Nutrient Management system of crops. The application of graded levels of major nutrients, with and without the combinations of micronutrients, vermin-compost and bio-fertilizers totaling fifteen treatments to sugarcane crop (Kanjana, 2007); and highest dose of 340, 80 and 140 kg N, P and K ha⁻¹ of N, P and K in combination with micronutrients of 37.5 kg ZnSO₄ and 100 kg FeSO₄ ha⁻¹), vermin-compost (5 t ha⁻¹) and Azophos (2.4 kg ha⁻¹) maintained higher levels of available nutrients at different stages of crop growth. There was great impact biocompost, farmyard manure (FYM) and press-mud cake (PMC) as well as integration of these organic fertilizers with inorganic N on the growth, yield, quality and leaf N content of sugarcane and soil nutrient status. Among the organics, FYM on equal N basis application increased the shoot population, number of millable canes and yield over others without remarkable change in sugar content of cane. Integrated application of either of the organics with inorganic exhibited better impact on the growth and yield characters and the 50:50 integration proved superior over others, closely followed by 33:67% ratio (Srivastava *et al.*, 2005). However the application of 25% less inorganic fertilizer to the recommended level of chemical fertilizer with press mud or FYM could be used to prevent nutrient depletion and maintain productivity as well (Bokhtiar, *et al.*, 2005). Organic manures applied to the crop markedly enhance the ratoon cane yield and juice quality and improve the physical conditions of the soil (Singh *et al.*, 2003). Organic fertilizer has

residual nitrogen (N) effect after the year of its application to land, as the decomposition of organic material usually takes more than a year (Lund & Doss 1980; Magdoff & Amadon 1980; Grolitz *et al.*, 1985; Werner *et al.*, 1985; Sommerfeldt *et al.*, 1988; Dilz *et al.*, 1990). Manures can supply nitrogen (N) beyond the year of its application, producing residual effects that are not fully expressed in short-term experiments (Schroder, 2005). Declining soil health and soaring prices of market purchased inputs, intensive cropping and imbalanced use of essential plant nutrients have rendered the alluvial soils of sub-tropics to be poor in organic carbon content and deterioration in physical properties (Speir *et al.*, 2004) lead to restricted growth and development of the crop. Keeping this in view the present study was planned to find agro-techniques which on one hand can ensure balanced supply of nutrients for the crop and on the other effectively restore the soil health and also save on the cost of inputs.

MATERIALS AND METHODS

The soil of experiment was clay loam in texture, EC (0.97 dSm⁻¹), slightly alkaline in reaction pH (7.7), calcareous (CaCO₃ 9.5%), low in organic matter (0.6%), total nitrogen content (0.05%) and available phosphorus (3.5 mg kg⁻¹), however high in exchangeable potassium (180 mg kg⁻¹). The experiment was conducted using randomized complete block design, the net plot size was kept 35m² and all the treatments were replicated three times.

Cultural practices: The land was prepared with disc plough to remove the hard pan of the soil and to achieve fine seed bed. The precision land leveling was maintained after bund making and developing plots according to the layout plan. The FYM, sugarcane press mud and bio fertilizer were thoroughly mixed with soil according to treatments i.e. one month before sowing of crop. The ridges were made at row spacing of 1 meter. The sets bearing 2-3 buds of sugarcane variety Thatta-10 were placed in the furrows (overlapping) and covered with thin layer of soil. A light irrigation was applied immediately after sowing. Later, the crop was irrigated at 15 days interval in winter months and at 8-10 days interval during summer months. Overall, 25 irrigations were applied during the growing season. Fertilizers were applied according to the experimental treatments. Nitrogen, phosphorous and potassium were applied in the form of Urea, DAP and SOP, respectively. Nitrogen was applied in three equal split doses i.e., at the time of sowing, after germination and completion of tillering. All phosphorus and potash were applied at the time of sowing. Two earthings were done both in plant as well as in ratoon crop. Cultural method of weed control was applied through interculturing. Stem borers were controlled by release of

Trichogramma chilonis (Lshii). Larsben was applied at the rate of 5 lit ha⁻¹ with 1st irrigation to control the termites. The data on cane yield, yield components and quality parameters were collected and statistically analyzed through “Statistix 8.1” computer software. The LSD value for mean comparison was calculated only if the general treatment *F* test was significant at a probability of ≤ 0.05 (Gomez & Gomez, 1984).

RESULTS AND DISCUSSION

Effect of inorganic and organic nutrients on agronomic traits of plant crop:

The results of the experiment revealed maximum germination (58.33%), tillers plant⁻¹ (5.83), plant height (266.78 cm), stem girth (2.63 cm), internodes plant⁻¹ (16.99), internode length (13.33 cm), millable canes (120.0 thousands ha⁻¹) and cane yield (118.33) of plant crop with the application of FYM @ 20 t ha⁻¹ and three-fourth of recommended NPK (169-84-126) and 49.33, 6.55, 278.89, 2.64, 16.22, 14.0, 123.66 and 119.60 respectively with application of press mud @ 20 t ha⁻¹ with three-fourth of recommended NPK (169-84.-126 kg ha⁻¹). The mean values of both the treatments were statistically non-significant. The results agree with the findings of Bokhtiar *et al.*, (2005) that cane height up to 3.19 m can be realized from press mud treatments followed by farmyard manure with cane and sugar yield of 199.14 t ha⁻¹ and 10.99 t ha⁻¹ respectively. The findings of Yadav *et al.*, (1993) also supports the results of this study that combined application of cow dung at 20 t ha⁻¹ with 50 kg N ha⁻¹ increased cane yield. Yield increase in sugarcane, have also been reported by Bevacqua & Mellano (1994); Hallmark *et al.*, (1995) with addition of organic amendments to soil. This study is fully supported by Rakkiyappan *et al.*, (2001) that highest cane and sugar yields could be achieved with the application of 75% NPK + pressmud, whereas, combined application of organic and chemical fertilizers significantly enhanced the yield over chemical fertilizers alone, Babaret *et al.*, (2011), also found that fifty percent bio-fertilizer + 50% recommended dose of chemical fertilizer increases the cane yield along with reduced environmental pollution. The results further showed that lower rates or individual application of inorganic or organic nutrient sources recorded less value of all agronomic traits. The minimum germination (49.66%), tillers plant⁻¹ (3.3), plant height (210.4 cm), stem girth (2.43- cm), internodes plant- 1 (12.5), internode length (7.8 cm), millable canes (44.0 thousands ha⁻¹) and cane yield (30.41) were found in untreated plots (Table 1).

Effect of inorganic and organic nutrients on physiological traits of plant crop:

The results of the experiment showed that integrated use of FYM or press mud with inorganic fertilizers significantly recorded higher values of all the physiological traits of plant crop. The maximum leaf area plant⁻¹ (5840 cm²), leaf area index (10.51), crop growth rate (10.35 gm⁻² day⁻¹) and dry matter (5542 gm⁻²) were found with the application of three-fourth of recommended fertilizer (169-84-126) in combination of PM @ 20 t ha⁻¹ and 5717, 10.29, 9.71 and 5206 respectively with the application of three-fourth of recommended fertilizer (169-84-126) in combination with FYM @ 20 t ha⁻¹. The mean values of these both treatments were statistically non-significant. However, under-dose applications of NPK fertilizer with organic sources significantly recorded less values of all the physiological traits of sugarcane plant crop. The results of the study are also supported by Nasir *et al.*, (2000) with opinion that higher growth rate of sugarcane was mainly due to the enhanced uptake of N, P and K. The minimum leaf area plant⁻¹ (3076 cm²), leaf area index (5.5), crop growth rate (6.13 gm⁻² day⁻¹) and dry matter (3286 gm⁻²) were noted in untreated plots (Table 2).

Effect of inorganic and organic nutrients on qualitative traits of plant crop:

The results of the experiment showed that application of three-fourth of the recommended NPK (169-84-126) + 20 t ha⁻¹ press mud was superior nutrient level for obtaining maximum brix (23.56%), pol (20.07%), purity (85.18%) and commercial cane sugar (14.67%) and 23.36, 19.76, 84.56 and 14.37 respectively with application of three-fourth of the recommended NPK (169-84-126) + 20 t ha⁻¹ FYM except fiber being higher (12.50%) in the unfertilized plots. The results are in agreement with the findings of Bokhtiar *et al.*, (2005) which reveals that press mud application with inorganic fertilizer increased brix, Pol and purity and thus showed better performance as compared to FYM or inorganic fertilizer alone. The minimum brix (20.76 %), pol (16.00 %), purity (77.04) and commercial cane sugar (10.86%) were recorded in control (Table 3).

Effect of inorganic and organic nutrients on NPK concentration and uptake of plant crop:

The results of the study showed maximum concentration of N (1.55%), P (0.44%) and K (1.73%) having N uptake (165.94 kg ha⁻¹), P uptake (48.64 kg ha⁻¹) and K uptake (163.01 kg ha⁻¹) with the application of three-fourth of recommended NPK (169-84-126) plus press mud @ 20 t ha⁻¹ and 1.54, 0.44, 1.21, 166.41, 47.77 and 130.18 respectively with the application of three fourth of recommended NPK (169-84-126) with FYM @ 20 t ha⁻¹. These results are supported by Bokhtiar *et al.*, (2001) that N, P and K was lower in sugarcane applied with inorganic

fertilizers than the crop added with press mud. Efficient storage and recycling of animal manure could contribute to improved P utilization, which in turn may reduce P fertilizer requirements (Haygarth & Jarvis, 1999). However, minimum NPK concentration (1.16, 0.19 and 1.21% respectively) and NPK uptake (61.23, 10.46 and 64.67 kg ha⁻¹ respectively) were observed in unfertilized plots (Table 4).

ECONOMIC ANALYSIS

Physical and revenue productivity:

The partial economic analysis sugarcane plant crop showed maximum revenue of Rs. 299000 from cane yield sale (119.6) with application of press mud applied at 20 t ha⁻¹+ three-fourth of the recommended inorganic NPK fertilizer (169-84-126) followed by revenue of Rs. 295825 from cane yield sale (118.33 t ha⁻¹) with application of farm yard manure applied at 20 t ha⁻¹ + three-fourth of the recommended inorganic NPK fertilizer (169-84-126) (Table 5).

Net returns:

The high net returns of Rs. 187935 ha⁻¹ were obtained with application of press mud applied at 20 t ha⁻¹+ three-fourth of the recommended inorganic NPK fertilizer (169-84.-126.). However, Rs. 179760 were obtained with the application of farm yard manure applied @ 20 t ha⁻¹+ three-fourth of the recommended inorganic NPK fertilizer (Table 5).

Cost benefit ratio:

The cost benefit ratio 6.36 and 5.48 were higher in the treatment where press mud and farm yard manure applied at 20 t ha⁻¹+ three-fourth of the recommended inorganic NPK fertilizer (169-84-126).

CONCLUSION

It is concluded that integrated nutrient management recorded 25% saving of inorganic fertilizers due to application of FYM and or press mud applied at 20 t ha⁻¹. Partial economic analysis showed higher revenue and net returns through integration of organic and inorganic nutrient sources. Integration of organic and inorganic nutrients should be practiced. This will not only enhance growth, yield, quality and nutrient uptake of sugarcane but also conserve agro-ecosystem for sustainable crop production.

Table 1. Effect of inorganic and organic nutrients on agronomic traits of plant crop

Treatments	Germination (%)	Tillers plant ⁻¹	Plant height (cm)	Stem girth (cm)	Millable canes (000) ha ⁻¹	Cane yield (t ha ⁻¹)
Control (untreated) 0-0-0 NPK kg ha ⁻¹	49.66g	3.7 gh	212.4 h	2.43 de	44.00 i	30.41 i
100% recommended (225-112-168 NPK kg ha ⁻¹)	55.00 bc	5.33 bc	260.7 bc	2.53 bc	96.30 cd	94.81 cd
Farm yard manure 10 t ha ⁻¹	50.00 fg	3.88 gh	213.1 h	2.44 cde	53.50 hi	42.70 hi
Farm yard manure 20 t ha ⁻¹	51.00 efg	4.22 fg	215.2 fgh	2.44 cde	62.00 gh	53.74 h
Press mudd 10 t ha ⁻¹	51.33 d-g	4.33 efg	222.44 f	2.46 cde	53.70 hi	42.43 hi
Press mudd 20 t ha ⁻¹	53.00 c-f	4.11 fg	222.33 fg	2.51 cd	68.73fg	54.00 h
Bio fertilizer F 5 t ha ⁻¹	52.00 c-g	3.44 h	210.00 h	2.42 e	45.33 i	34.24 i
Bio fertilizer 10 t ha ⁻¹	50.66 fg	4.10 fg	217.00 fgh	2.46 cde	48.76 hi	39.01 i
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	54.33 bcd	5.22 bcd	263.78 b	2.53 bc	107.17 bc	105.80 bc
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	58.33 a	5.83 b	266.78 b	2.63 a	120.00 a	118.33 a
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	57.33 ab	5.83 b	267.00 b	2.61 ab	106.30 bc	106.15 bc
25% less (168.75-84.0-126.0 NPK kg) + PM 20 t ha ⁻¹	49.33 g	6.55 a	278.89 a	2.64 a	123.66 a	119.60 a
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	54.00 cde	4.55 ef	250.00 d	2.44 cde	87.70 de	85.59 def
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	54.00 cde	4.88 cde	255.89 cd	2.45 cde	90.60 de	88.11 fg
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	50.66 fg	4.66 def	232.89 e	2.44 cde	80.50 ef	73.04 fg
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	51.66 d-g	4.88 cde	234.44 e	2.47 cde	82.60 e	76.00 efg
50% less (112.5-56.0-84.0 NPK kg) + PM 10 t ha ⁻¹	52.00 c-g	5.22 bcd	235.00 e	2.47 cde	84.00 de	74.33 fg
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 20 t ha ⁻¹	49.66 g	5.55 b	237.22 e	2.49 cde	86.90 de	76.85 efg
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	52.00 c-g	3.33 h	215.00 gh	2.48 cde	81.33 ef	70.15 g
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	51.33 d-g	4.21 fg	214.89 h	2.44 cde	82.00 ef	72.30 g
S.E	1.6228	0.3167	3.6511	0.0466	6.687	6.164
LSD (0.5%)	3.2851	0.6412	7.3913	0.0943	13.539	12.479

PM= Press mud, FYM= Farm yard manure, B.F= Biofertilizer

Table 2. Physiological parameters as affected by organic and in organic plant nutrient management in sugarcane plant crop.

Treatments	Leaf Area plant ⁻¹ (cm ²)	Leaf Area Index	Crop Growth Rate (g m ² day ⁻¹)	Dry matter (gm ²)
Control (untreated) 0-0-0 NPK kg ha ⁻¹	3076 i	5.537 h	6.133 l	3286 k
100% recommended (225-112-168 NPK kg ha ⁻¹)	5481 a	9.870 a	9.110 c	4880 c
Farm yard manure 10 t ha ⁻¹	3270 hi	5.890 gh	6.327 kl	3390 jk
Farm yard manure 20 t ha ⁻¹	3424 fghi	6.167 efgh	6.593 jk	3543 ijk
Press mudd 10 t ha ⁻¹	4478 bc	8.060 b	6.733 ij	3607 hijk
Press mudd 20 t ha ⁻¹	3632 efgh	6.540 defg	7.023 hi	3695 ghij
Bio fertilizer F 5 t ha ⁻¹	3360 ghi	6.050 fgh	6.217 kl	3331 k
Bio fertilizer 10 t ha ⁻¹	3537 efgh	6.370defg	6.563 jk	3516 ijk
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	5623 a	10.12 a	9.330 c	4999 bc
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	5717 a	10.29 a	9.717 b	5206 b
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	5755 a	10.36 a	9.860 b	5282 ab
25% less (168.75-84.0-126.0 NPK kg) + PM 20 t ha ⁻¹	5840 a	10.51 a	10.35 a	5542 a
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	4460 bc	8.030 b	7.467 efg	4000 defg
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	4573 b	8.230 b	7.897 d	4197 de
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	3760 ef	6.770 def	7.263 fgh	3891 efgh
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	3837 de	6.910 cde	7.453 efg	4292 d
50% less 112.5-56.0-84.0 NPK kg + PM 10 t ha ⁻¹	4192 cd	7.550 bc	7.550 def	4035 def
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 20 t ha ⁻¹	3919 de	7.050 cd	7.690 de	4121 def
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	3715 efg	6.990 cd	7.113 gh	3810 fghi
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	3813 e	6.860 cde	7.460 efg	3996 defg
S.E	120.4	0.2373	0.1238	101.5
LSD (5%)	344.4	0.6795	0.3545	290.6

Table 3. Quality traits of sugarcane crop as affected by organic and inorganic plant nutrients in sugarcane plant crop.

Treatments	Fiber (%)	Brix (%)	Pol (%)	Purity (%)	CCS (%)
Control (untreated) 0-0-0 NPK kg ha ⁻¹	12.50 a	20.76 q	16.00 r	77.04 mn	10.86 r
100% recommended (225-112-168 NPK kg ha ⁻¹)	12.24 j	22.90 e	19.26 e	84.10 d	13.97 e
Farm yard manure 10 t ha ⁻¹	12.38 ef	21.00 o	16.20 p	77.15 lm	11.02 p
Farm yard manure 20 t ha ⁻¹	12.39 ef	21.16 m	16.35 o	77.27 klm	11.13 o
Press mudd 10 t ha ⁻¹	12.44 bc	21.36 l	16.53 n	77.37 kl	11.26 n
Press mudd 20 t ha ⁻¹	12.44 bc	21.66 k	16.79 m	77.51 k	11.46 m
Bio fertilizer F 5 t ha ⁻¹	12.44 bc	20.86 p	16.09 q	77.13 lm	10.94 q
Bio fertilizer 10 t ha ⁻¹	12.40 cde	21.06 n	16.25 p	76.79 n	11.05 p
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	12.37 ef	23.16 d	19.56 d	84.45 c	14.21 d
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	12.33 gh	23.36 c	19.76 c	84.56 bc	14.37 c
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	12.27 ij	23.46 b	19.91 b	84.86 ab	14.52 b
25% less (168.75-84.0-126.0 NPK kg) + PM 20 t ha ⁻¹	12.25 j	23.56 a	20.07 a	85.18 a	14.67 a
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	12.38 ef	22.76 f	18.94 g	83.20 e	13.62 g
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	12.36 fg	22.86 e	19.06 f	83.37 e	13.73 f
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	12.43 bcd	22.56 h	18.17 j	80.54 h	12.77 j
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	12.44 bc	22.66 g	18.26 i	80.57 h	12.87 j
50% less 112.5-56.0-84.0 NPK kg + PM 10 t ha ⁻¹	12.32 gh	22.76 f	18.73 h	82.29 g	13.37 i
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 20 t ha ⁻¹	12.29 hi	22.86 e	18.90 g	82.66 f	13.53 h
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	12.45 b	21.76 j	17.22 l	79.10 j	11.94 l
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	12.40 de	22.03 i	17.52 k	79.52 i	12.19 k
S.E	0.0191	0.0186	0.0283	0.1595	0.0346
LSD (5%)	0.0387	0.0376	0.0572	0.3229	0.0700

Table 4. NPK concentration and uptake as affected by inorganic and organic fertilizers in sugarcane plant crop

Treatments	N (%)	P (%)	K (%)	N uptake (kg ha ⁻¹)	P uptake (kg ha ⁻¹)	K uptake (kg ha ⁻¹)
Control (untreated) 0-0-0 NPK kg ha ⁻¹	1.16 ij	0.33 g	1.21 q	88.76 j	24.82 i	84.8 2 k
100% recommended (225-112-168 NPK kg ha ⁻¹)	1.44 ef	0.42 d	1.58 f	140.85 c	41.60 de	155.19 b
Farm yard manure 10 t ha ⁻¹	1.51 b	0.42 d	1.46 h	102.23 i	28.52 gh	99.29 i
Farm yard manure 20 t ha ⁻¹	1.54 a	0.41 d	1.66 b	104.32 hi	28.29 gh	112.90 g
Press mudd 10 t ha ⁻¹	1.42 g	0.42 d	1.48 g	100.21 i	29.99 gh	122.09 f
Press mudd 20 t ha ⁻¹	1.45 de	0.37 f	1.63 c	103.27 i	26.58 hi	116.09 g
Bio fertilizer F 5 t ha ⁻¹	1.34 i	0.40 e	1.61 d	101.09 i	26.62 hi	106.65 h
Bio fertilizer 10 t ha ⁻¹	1.47 c	0.44 bc	1.39 k	98.95 i	29.59 gh	93.37 j
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	1.50 b	0.41 d	1.43 i	157.55 b	43.64 cd	149.54 c
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	1.54 a	0.45 a	1.30 n	166.41 a	47.77 ab	130.18 e
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	1.46 d	0.42 cd	1.41 j	153.97 b	45.17 bc	148.70 c
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + PM 20 t ha ⁻¹	1.55 a	0.44 ab	1.73 a	165.94 a	48.64 a	163.01 a
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	1.42 g	0.42 d	1.49 g	130.06 d	38.86 ef	136.47 d
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	1.43 ef	0.44 ab	1.58 f	132.33 d	42.18c-e	145.53 c
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	1.38 h	0.42 cd	1.60 e	114.60 fg	35.56 f	132.86 de
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	1.43 fg	0.42 d	1.63 c	120.05 e	35.73 f	136.84 d
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	1.42 fg	0.44 ab	1.38 l	126.97 d	29.81 gh	122.82 f
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 20 t ha ⁻¹	1.34 i	0.34 g	1.41 j	119.59 ef	30.73 g	125.84 f
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	1.22 j	0.33 g	1.37 m	91.68 j	28.96 gh	102.96 hi
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	1.43 ef	0.36 f	1.39 k	109.62 gh	27.84 ghi	106.06 h
S.E	7.574 E	7.556 E	4.806 E	2.6845	1.6861	2.0119
LSD (5%)	0.015	0.015	9.730	0.1382	0.0847	0.414

Table 5 Partial economic analysis of sugarcane plant crop under the influence of organic and inorganic nutrient sources.

Treatments	Physical productivity (t ha ⁻¹)	Revenue productivity (Rs)	Cost of production (Rs)	Net return (Rs)	Cost benefit ratio
Control (untreated) 0-0-0 NPK kg ha ⁻¹	30.41	76025	0	-	-
100% recommended (225-112-168 NPK kg ha ⁻¹)	94.81	237025	33386	127614	4.82
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) +FYM 10 t ha ⁻¹	105.8	264500	32540	155935	5.79
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) +FYM 20 t ha ⁻¹	118.33	295825	40040	179760	5.48
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	106.15	265375	30040	159310	6.3
25% less (168.75-84.0-126.0 NPK kg) + PM 20 t ha ⁻¹	119.6	299000	35040	187935	6.36
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	85.59	213975	35040	102910	3.93
25% less (168.75-84.0-126.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	88.11	220275	45040	99210	3.2
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 10 t ha ⁻¹	73.04	182600	24194	82381	4.4
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + FYM 20 t ha ⁻¹	76.0	190000	31694	82281	3.59
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 10 t ha ⁻¹	74.33	185825	21694	79194	5.06
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + PM 20 t ha ⁻¹	76.85	192125	26694	89406	4.34
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 5 t ha ⁻¹	70.15	175375	26694	72656	3.72
50% less (112.5-56.0-84.0 NPK kg ha ⁻¹) + BF 10 t ha ⁻¹	72.30	180750	36694	68031	2.85

Cost of fertilizer: Urea Rs.14/kg, TSP Rs.40/kg and SOP Rs.50/kg.

Cost of Manures: Press mud Rs. 500/t, FYM Rs. 875/t and Biofertilizer Rs.4000/t

Cost of produce: Cane Rs.2500/t

References

- Babar, L.K., T. Iftikhar, H.N. Khan and A.H. Makhdam. 2011. Agronomic trials on sugarcane crop under Faisalabad conditions, Pakistan. *Pak. J. Bot.*, 43(2): 929-935.
- Bhander, P.K., M.S.U. Bhuiya and M.A. Salam. 1998. Effect of *Sesbania rostrata* biomass and nitrogen fertilizer on the yield and yield attributes of transplant Amain rice. *Progressive Agric.*, 9: 89-93.
- Bhattacharya, P. and D. Gehlot. 2003. Current status of organic farming at international and national level. *Agrobios News Letter*, 4: 7-9.
- Bevacqua, R.F. and V.J. Mellano. 1994. Cumulative effects of sludge compost on crop yields and soil properties. *Commun. Soil Sci. Plant Anal.*, 25: pp. 395.
- Bokhtiar, H., R.C. Rehman and A.B.M.M. Rahman. 2005. Efficacies of some organic fertilizers on sustainable sugarcane production in old Himalayan piedmont plain soil of Bangladesh. *Pak. Sug. J.*, 20(1): 2-5.
- Bokhtiar, S.M. and K. Sakurai. 2005. Integrated use of organic manure and chemical fertilizer on growth, yield and quality of sugarcane in high Ganges river flood plain soils of Bangladesh. *Commun. Soil Sci. Plant Anal.*, 36: 1823-1837.
- Bokhtiar, S.M., G.C. Paul, M.A. Rashid and A.B.M. Rahman. 2001. Effect of pressmud and organic nitrogen on soil fertility and yield of sugarcane grown in high Ganges river flood plain soils of Bangladesh. *Indian Sugar. L.*, 1: 235-240.
- Dang, Y.P., K.S. Verma and B.S. Pannu. 1995. Need for potassium fertilization in sugarcane. *Indian Sugar. XLV*: 229-235.
- Dutta, S., R. Pal, A. Chakeraborty and K. Chakrabarti. 2003. Influence of integrated plant nutrient supply system on soil quality restoration in a red and laterite soil. *Archives of Agron. and Soil Sci.*, 49: 631-637.
- Dilz, K.K., J. Postmus and W.H. Prins. 1990. Residual effect of long term applications of farmyard manure to silage maize. *Fertilizer Res.*, 26: 249-252.
- Ghugare, R.V., S.S. Magar and S.Y. Daftardar. 1988. Effect of distillery effluent (spentwash) with dilution on growth and yield parameters of adseli sugarcane (Co 740). Paper in National Seminar on Sugar Factory and Allied Industrial Wastes-A new focus: 1-3.
- Gomez, K.A. and A.A. Gomez. 1984. Statistical Procedure for Agricultural Research, (2 eds.), Wiley, New York, USA. pp. 680.

- Gorlitz, H., V. Herrmann and R. Jauert. 1985. Ertrag und Nährstoffnutzung nach ein- und mehrjährigen hohen Gullegaben zu Silomais sowie ihre Nachwirkung auf sandigen Böden. *Arch. Acker- Pflanzenb. Bodenk. Berlin*, 29: 55-60.
- Hallmark, W.B., S.E. Feagley, G.A. Breitenbeck, L.P. Brown, X. Wan and G.L. Hawkins. 1995. Use of composted municipal waste in sugarcane production. *Louisiana Agric.*, 38: 15-16.
- Haygarth, P.M. and S.C. Jarvis. 1999. Transfer of phosphorus from agricultural soils. *Adva. Agron.*, 66: 195-249. Kumaraswamy, K., R.V. Reddy and K. Babu. 1998 Cumulative effect of continuous cropping and manuring of sugarcane on organic matter and NPK status of the soil. *J. Indian Soc. Soil Sci.*, (46): 47-49.
- Khan, I.A., S. Bibi, S. Yasmin, A. Khatri, N. Semma and S.A. Abro. 2012. Correlation studies of agronomic traits of higher sugar yield in sugarcane. *Pak. J. Bot.*, 44(3): 969-971.
- Kanjana, D., G. Pitchai and J. Saravanan. 2007. Effect of organic, inorganic and biofertilizer on soil nutrient availability in sugarcane variety (CO 86032) cultivation at Theni District of Tamil Nadu. *Indian Sugar*, 56(11): 15-22.
- Lund, Z.F. and B.D. Doss. 1980. Residual effects of dairy cattle manure on plant growth and soil properties. *Agron. J.*, 72: 123-130.
- Magdoff, F.R. and J.F. Amadon. 1980. Yield trends and soil chemical changes resulting from N and manure application to continuous corn. *Agron. J.*, 72: 161-164.
- Mumtaz, A.S., Dur-e-Nayab, M.J. Iqbal and Z.K. Shinwari. 2011. Probing genetic diversity to characterize red rot resistance in sugarcane. *Pak. J. Bot.*, 43(5): 2513-2517
- Murphy, J. and J.P. Riley. 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal. Chim. Acta.*, 27: 31-36.
- Nasir, M.N., R. H. Qureshi, M. Aslam and J. Akhtar. 2000. Comparative studies on two selected sugarcane lines for differences in physiological traits. *Pak. Sugar*, 6: 52-61.
- Nawab, K., Amanullah, P. Shah, Abdur-Rab, M. Arif, M.A. Khan, A. Mateen and F. Munsif. 2011. Impact of integrated nutrient management on growth and grain yield of wheat under irrigated cropping system. *Pak. J. Bot.*, 43(4): 1943-1947.
- Palaniappan, S.P. and K. Annadurai. 1999. *Organic Farming Theory and Practice*, Jodhpur India, Scientific Publishers. pp. 53-73.
- Rakkiyappan, P., S. Thangavelu, R. Malathi and R. Radhamani. 2001. Effect of biocompost and enriched pressmud on sugarcane yield and quality, 3(3): 92-96.
- Radford, P. J. 1967. Growth analysis of formulae, their use and abuse. *Crop Sci.*, 7: 171-175.

- Saviozzi, A., P. Bufalino, R. Levi Minzi and R. Riffaldi. 2002. Biochemical activities in a degraded soil restored by two amendments: a laboratory study. *Biology and Fertility of Soils*, 35: 96-119.
- Singh, S.N., A.K. Singh, S.C. Singh, M.L. Sharma and R. Kumar. 2011. Enhancing sugarcane (*Saccharum* spp. Hybrid) productivity by integrating organic, inorganic and biological sources of N in sub-tropical India. *Indian Journal of Sugar Technology*, 26(1): 14-15.
- Srivastava, T.K., M. Lal, K.P. Singh, A. Suman and P. Kumar. 2009. Enhancing soil health and sugarcane productivity in a plant-ratoon system through organic nutrition modules in subtropics. *Indian J. Agricultural Sciences*, 79(5): 346-350.
- Srivastava, P.N., D.N. Singh and S.B. Singh. 2005. Integrated nutrient management in sugarcane: performance of FYM, Biocompost and pressmud cake with inorganic N on growth and quality indices. *Cooperative Sugar*, 36(12):993-998.
- Sommerfeldt, T.G., C. Chang and T. Entz. 1988. Long-term annual manure applications increase soil organic matter and nitrogen, and decrease carbon to nitrogen ratio. *Soil Sci.Soc. America. J.*, 52: 1668-1672.
- Schroder, J.J. 2005. Revisiting the agronomic benefits of manure: a correct assessment and exploitation of its fertilizer value spares the environment. *Bio resource Tech.*, 96(2): 253-261.
- Singh, I.S., R. Singh, M.P. Yadav, R.R. Singh and S.B. Singh. 2003. Ratooning behaviour of new promising varieties of sugarcane under Bhat soil conditions. *Indian Sugar*, LIII(5): 333-336.
- Spier, T.W., J. Horswell, R.G. McLaren, G. Fietje and A.P. Van Schalk. 2004. Composted biosolids enhance fertility of sandy loam soil under dairy pasture. *Biology and Fertility of Soils*, 40: 349-358.
- Werner, W., H.W. Scherer and D. Drescher. 1985. Untersuchungen über den Einfluss langjähriger Gülledüngung auf N-Fractionen und N-Nachlieferung des Bodens. *Zeitschr. Acker-Pflanzenb.*, 155: 137-144.
- Yadav, R.L. 1993. Agronomy of sugarcane: Principle and Practice. 1st Eds. International Book Distributing Co. Lucknow, India.